



HAL
open science

On some constants related to the harmonic zeta function

Marc-Antoine Coppo, Bernard Candelpergher

► **To cite this version:**

Marc-Antoine Coppo, Bernard Candelpergher. On some constants related to the harmonic zeta function. 2024. hal-03602568v2

HAL Id: hal-03602568

<https://univ-cotedazur.hal.science/hal-03602568v2>

Preprint submitted on 19 May 2024 (v2), last revised 8 Oct 2024 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

On some constants related to the harmonic zeta function

Marc-Antoine Coppo* and Bernard Candelpergher

Université Côte d'Azur, CNRS, LJAD (UMR 7351), Nice, France

Abstract We study a range of constants that are closely linked to the harmonic zeta function ζ_H . In addition, by means of a new representation of ζ_H , we derive possibly new integral formulas for $\zeta(n)$ and $\zeta'(n)$ and new evaluations for a variety of integrals of the same type as those previously considered by Glasser, Manna, and Oloa.

Keywords Harmonic zeta function; Stieltjes constants; zeta values; Glasser-Manna-Oloa integrals, Ramanujan summation of series.

1 Introduction

Let ζ_H be the harmonic zeta function defined for $\operatorname{Re}(s) > 1$ by

$$\zeta_H(s) := \sum_{n=1}^{\infty} \frac{H_n}{n^s},$$

where, for all $n \geq 1$,

$$H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}$$

are the classical harmonic numbers. Four decades ago, Apostol and Vu [1] showed that this function could be continued as a meromorphic function with a double pole at $s = 1$, and an infinite number of simple poles at $s = 0$ and $s = 1 - 2k$ for each integer $k \geq 1$. The Laurent expansion of the harmonic zeta function ζ_H around its double pole $s = 1$ can be written as

$$\zeta_H(s) = \frac{1}{(s-1)^2} + \frac{\gamma}{s-1} + \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \gamma_H^{(k)} (s-1)^k \quad (0 < |s-1| < 1),$$

*Corresponding author. *Email address:* coppo@unice.fr

where $\gamma = -\Gamma'(1)$ is Euler's constant, and the coefficients $\gamma_H^{(k)}$ are called *harmonic Stieltjes constants* by analogy with the classical Stieltjes constants γ_k . A common definition of these constants is

$$\gamma_k = \lim_{s \rightarrow 1} \left\{ (-1)^k \zeta^{(k)}(s) - \frac{k!}{(s-1)^{k+1}} \right\} \quad (k \geq 0),$$

where $\zeta^{(k)}(s)$ is the k th derivative of the Riemann zeta function. In section 3, we give a new evaluation of the harmonic Stieltjes constants $\gamma_H^{(k)}$ (see Corollary 1) which arises from a new expression of $\zeta_H(s)$. From now on and throughout the text, we will use the lighter notation $\gamma^{(k)}$ in place of $\gamma_H^{(k)}$. An explicit expression of $\gamma^{(0)}$ is given by the following formula [9, Equation (6)]:

$$\gamma^{(0)} = \frac{1}{2}\gamma^2 + \frac{1}{2}\zeta(2) = \frac{1}{2}\Gamma''(1). \quad (1)$$

It is worth noting that this nice identity extends to the case of the height 1 multiple zeta functions $\zeta(s, 1, \dots, 1)$ [17, Corollary 4.1]. Regarding the classical Stieltjes constants, we recall the well-known asymptotic formula

$$\gamma_k = \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N \frac{(\log n)^k}{n} - \frac{(\log N)^{k+1}}{k+1} \right\} \quad (k \geq 0).$$

From our point of view, it is important to point out that γ_k is nothing else than the \mathcal{R} -sum (i.e. the sum of the series in the sense of Ramanujan's summation method) of the divergent series $\sum_{n \geq 1} \frac{(\log n)^k}{n}$ [8, p. 67]. The existence of a similar asymptotic formula for $\gamma^{(k)}$:

$$\gamma^{(k)} = \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N \frac{H_n (\log n)^k}{n} - \frac{(\log N)^{k+2}}{k+2} - \gamma \frac{(\log N)^{k+1}}{k+1} \right\} \quad (k \geq 0),$$

which results from the extension of a formula of Briggs and Buschman [3, 7], strongly suggests that the constant $\gamma^{(k)}$ is closely linked to the \mathcal{R} -sum of the divergent series $\sum_{n \geq 1} \frac{H_n (\log n)^k}{n}$. This is indeed the case: such a relation exists although it is more intricate than in the classical case. We have already encountered in earlier studies [8, Equation (3.23)], [9, Equation (4)] the following relation :

$$r_0 = \gamma^{(0)} + \tau_1 + \gamma_1 - \zeta(2) = \frac{1}{2}\gamma^2 - \frac{1}{2}\zeta(2) + \gamma_1 + \tau_1, \quad (2)$$

where

$$r_0 := \sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n} \quad \text{and} \quad \tau_1 = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\zeta(n+1)}{n} = - \sum_{n=2}^{\infty} \zeta'(n).$$

We will see in the next section to what extent our formula (2) can be generalized (see Proposition 1). In the subsequent section, the new representation of ζ_H provided by Theorem 1, which is an improvement of a formula due to Boyadzhiev, Gadiyar and Padma [6], allows us to derive new integral formulas for $\zeta(n)$ and $\zeta'(n)$ (see Corollary 2), as well as new evaluations for a variety of integrals of the same type as those previously considered by Glasser and Manna [14], and Oloa [16] (see Corollaries 4–6).

2 How the constants $\gamma^{(k)}$ and $\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n (\log n)^k}{n}$ are linked together

The forthcoming Proposition 1 successfully generalizes the previous relation (2).

Proposition 1. Let k be a natural number. If r_k and τ_k are defined by

$$r_k := \sum_{n \geq 1}^{\mathcal{R}} \frac{H_n (\log n)^k}{n} \quad (k \geq 0),$$

and

$$\tau_k := \sum_{n=1}^{\infty} (-1)^{n+k} \frac{\zeta(n+1)}{n^k} \quad (k \geq 1),$$

then

$$r_0 = \gamma^{(0)} + \tau_1 + \gamma_1 - \zeta(2),$$

and the following relations hold true for all positive values of k :

$$r_{2k-1} = \gamma^{(2k-1)} + (2k-1)! \tau_{2k} + \frac{\gamma_{2k}}{2k} + \sum_{n=1}^k \frac{(2k-1)!}{(2k-2n)!} \times \frac{(2^{2n-1} - 1)}{2^{2n-2}} \gamma_{2k-2n} \zeta(2n), \quad (3)$$

and

$$r_{2k} = \gamma^{(2k)} + (2k)! \tau_{2k+1} + \frac{\gamma_{2k+1}}{2k+1} + \sum_{n=1}^k \frac{(2k)!}{(2k-2n+1)!} \times \frac{(2^{2n-1} - 1)}{2^{2n-2}} \gamma_{2k-2n+1} \zeta(2n) - (2k)! \frac{(2^{2k+1} - 1)}{2^{2k}} \zeta(2k+2). \quad (4)$$

In particular,

$$\begin{aligned}
r_1 &= \gamma^{(1)} + \tau_2 + \frac{1}{2}\gamma_2 + \gamma\zeta(2), \\
r_2 &= \gamma^{(2)} + 2\tau_3 + \frac{1}{3}\gamma_3 + 2\gamma_1\zeta(2) - \frac{7}{2}\zeta(4), \\
r_3 &= \gamma^{(3)} + 6\tau_4 + \frac{1}{4}\gamma_4 + 3\gamma_2\zeta(2) + \frac{21}{2}\gamma\zeta(4), \\
r_4 &= \gamma^{(4)} + 24\tau_5 + \frac{1}{5}\gamma_5 + 4\gamma_3\zeta(2) + 42\gamma_1\zeta(4) - \frac{93}{2}\zeta(6).
\end{aligned}$$

Example 1. Numerical evaluations of $\gamma^{(k)}$, r_k and τ_k for $k \leq 4$ are given below.

$\gamma^{(0)} = 0.989055 \dots$	$r_0 = 0.529052 \dots$	τ_0 is undefined
$\gamma^{(1)} = 0.400761 \dots$	$r_1 = -0.078850 \dots$	$\tau_1 = 1.257746 \dots$
$\gamma^{(2)} = 0.971304 \dots$	$r_2 = -0.008095 \dots$	$\tau_2 = -1.424248 \dots$
$\gamma^{(3)} = 2.972584 \dots$	$r_3 = 0.003653 \dots$	$\tau_3 = 1.523800 \dots$
$\gamma^{(4)} = 11.940501 \dots$	$r_4 = 0.002816 \dots$	$\tau_4 = -1.580233 \dots$

Remark 1. Although not used afterwards, we also mention another expression of the constant τ_k which can be easily deduced from its definition:

$$\tau_k = (-1)^k \sum_{n=1}^{\infty} \frac{1}{n} \text{Li}_k \left(-\frac{1}{n} \right) \quad (k \geq 1),$$

where Li_k denotes the k th polylogarithm. In particular, for $k = 1$, the formula

$$\tau_1 = \sum_{n=1}^{\infty} \frac{1}{n} \log \left(1 + \frac{1}{n} \right)$$

is regained. Note that the definition of τ_k given here differs from that given in [9] when $k > 1$.

Proof of Proposition 1. The key formula to derive the general relations (3) and (4) is the decomposition of $\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^s}$ given by [9, Theorem 1] which is reproduced below:

$$\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^s} = \frac{\pi}{\sin(\pi s)} \zeta(s) + \int_0^1 \frac{\psi(x+1) + \gamma}{x^s} dx + \zeta_H(s), \quad (5)$$

where $\psi(x) = \frac{d}{dx} \log \Gamma(x)$ is the digamma function. This formula applies to all complex numbers s such that $\text{Re}(s) < 2$ and $s \neq 1, 0, 1 - 2k$ for each integer $k > 0$. Fortunately, the Laurent expansion of each component in (5) can be written explicitly.

a) The expansion of $\frac{\pi}{\sin(\pi s)} \zeta(s)$ at $s = 1$ can be obtained as follows: first, we write the successive equations:

$$\begin{aligned} \frac{-\pi}{\sin(\pi s)} &= \frac{\pi}{\sin(\pi(s-1))} = \frac{\exp(i\pi(s-1))}{s-1} \frac{2i\pi(s-1)}{\exp(2i\pi(s-1)) - 1} \\ &= \frac{1}{s-1} \left(\sum_{k \geq 0} i^k \pi^k \frac{1}{k!} (s-1)^k \right) \left(\sum_{k \geq 0} i^k (2\pi)^k \frac{B_k}{k!} (s-1)^k \right) \\ &= \frac{1}{s-1} + \sum_{k \geq 1} (-1)^k \left(\sum_{j=0}^{2k} \binom{2k}{j} 2^j B_j \right) \frac{\pi^{2k}}{(2k)!} (s-1)^{2k-1}, \end{aligned}$$

where B_j are the Bernoulli numbers. Euler's identity:

$$\zeta(2k) = (-1)^{k+1} 2^{2k-1} B_{2k} \frac{\pi^{2k}}{(2k)!} \quad (k \geq 1),$$

then allows us to rewrite this expansion as follows:

$$\frac{-\pi}{\sin(\pi s)} = \frac{1}{s-1} - \sum_{k \geq 1} \frac{2^{1-2k}}{B_{2k}} \sum_{j=0}^{2k} \binom{2k}{j} 2^j B_j \zeta(2k) (s-1)^{2k-1}.$$

Moreover, the latter expression can be simplified thanks to the identity

$$\sum_{j=0}^k \binom{k}{j} 2^j B_j = 2^k B_k \left(\frac{1}{2}\right) = 2(1 - 2^{k-1}) B_k \quad (k \geq 2).$$

Hence, the expansion of $\frac{\pi}{\sin(\pi s)}$ at $s = 1$ is given by

$$\frac{\pi}{\sin(\pi s)} = -\frac{1}{s-1} - \sum_{k=1}^{\infty} \frac{2^{2k-1} - 1}{2^{2k-2}} \zeta(2k) (s-1)^{2k-1}.$$

On the other hand, the expansion of $\zeta(s)$ at $s = 1$ is

$$\zeta(s) = \frac{1}{s-1} + \gamma + \sum_{k=1}^{\infty} \frac{(-1)^k}{k!} \gamma_k (s-1)^k,$$

where γ_k are the Stieltjes constants. The expansion of $\frac{\pi}{\sin(\pi s)} \zeta(s)$ is then obtained by Cauchy product. We have

$$\begin{aligned} \frac{\pi}{\sin(\pi s)} \zeta(s) &= -\frac{1}{(s-1)^2} - \frac{\gamma}{(s-1)} + \gamma_1 - \zeta(2) \\ &\quad - \left(\frac{1}{2} \gamma_2 + \gamma \zeta(2) \right) (s-1) + \left(\frac{1}{6} \gamma_3 + \gamma_1 \zeta(2) - \frac{7}{4} \zeta(4) \right) (s-1)^2 \\ &\quad - \left(\frac{1}{24} \gamma_4 + \frac{1}{2} \gamma_2 \zeta(2) + \frac{7}{4} \gamma \zeta(4) \right) (s-1)^3 \\ &\quad + \left(\frac{1}{120} \gamma_5 + \frac{1}{6} \gamma_3 \zeta(2) + \frac{7}{4} \gamma_1 \zeta(4) - \frac{31}{16} \zeta(6) \right) (s-1)^4 - \dots \quad (6) \end{aligned}$$

In (6), the coefficient of $(s-1)^{2k}$ and $(s-1)^{2k-1}$ are respectively

$$\frac{\gamma_{2k+1}}{(2k+1)!} + \sum_{j=1}^k \frac{\gamma_{2k-2j+1}}{(2k-2j+1)!} \times \frac{(2^{2j-1}-1)}{2^{2j-2}} \zeta(2j) - \frac{(2^{2k+1}-1)}{2^{2k}} \zeta(2k+2),$$

and

$$-\frac{\gamma_{2k}}{(2k)!} - \sum_{j=1}^k \frac{\gamma_{2k-2j}}{(2k-2j)!} \times \frac{(2^{2j-1}-1)}{2^{2j-2}} \zeta(2j).$$

b) The well-known Taylor series expansion of the digamma function:

$$\psi(x+1) + \gamma = \sum_{n=1}^{\infty} (-1)^{n+1} \zeta(n+1) x^n \quad (|x| < 1),$$

allows us write

$$\int_0^1 \frac{\psi(x+1) + \gamma}{x^s} dx = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\zeta(n+1)}{n-(s-1)}.$$

This leads to the following expansion:

$$\begin{aligned} \int_0^1 \frac{\psi(x+1) + \gamma}{x^s} dx &= \sum_{k=0}^{\infty} \left(\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\zeta(n+1)}{n^{k+1}} \right) (s-1)^k \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \times k! \tau_{k+1} (s-1)^k \quad (|s-1| < 1). \end{aligned} \quad (7)$$

c) To obtain the expansion of $\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^s}$ at $s=1$, we proceed as follows: first we write the identities

$$\frac{H_n}{n^s} = \frac{H_n}{n} e^{-(s-1)\log n} = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \times \frac{H_n (\log n)^k}{n} (s-1)^k,$$

so that

$$\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^s} = \sum_{n \geq 1}^{\mathcal{R}} \left(\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \times \frac{H_n (\log n)^k}{n} (s-1)^k \right).$$

Moreover, [8, Theorem 9] allows us interchange $\sum_{n \geq 1}^{\mathcal{R}}$ and $\sum_{k=0}^{\infty}$. This leads to the expansion:

$$\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^s} = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \left(\sum_{n \geq 1}^{\mathcal{R}} \frac{H_n (\log n)^k}{n} \right) (s-1)^k = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} r_k (s-1)^k. \quad (8)$$

The desired formulas (3) and (4) can then be easily obtained by combining the previous expansions (6), (7), and (8). □

3 A new expression of $\zeta_H(s)$ and its implications

In this section, we provide a new expression of the harmonic zeta function (Theorem 1) which allows us to give a new evaluation of the harmonic Stieltjes constants (Corollary 1). Thanks to this representation of ζ_H , we also derive new integral formulas for $\zeta(n)$ and $\zeta'(n)$ (Corollary 2), and new evaluations for Glasser-Manna-Oloa type integrals (Corollaries 4–6).

Theorem 1. *For all complex numbers s in $\mathbb{C} \setminus \mathbb{Z}$, we have*

$$\zeta_H(s) = \pi \cot(\pi s) \zeta(s) + \zeta(s+1) + \Gamma(1-s)\Phi(s) \quad (9)$$

with

$$\Phi(s) := -\frac{1}{2i\pi} \int_{-\pi}^{\pi} x \left(\text{Log}(1 + e^{ix}) \right)^{s-1} dx, \quad (10)$$

where Log denotes the principal value of the complex logarithm.

Proof. Let

$$z = \text{Log}(1 + e^{ix}) = \log\left(2 \cos\left(\frac{x}{2}\right)\right) + i \frac{x}{2}, \quad -\pi < x < \pi.$$

When x varies from $-\pi$ to π , the variable z traverses the path L defined by the parametric equations $\text{Re } z = \log(2 \cos(x/2))$ and $\text{Im } z = x/2$. This path extends from $-\infty$ below the line $\text{Im } z = 0$, passes through the point $(\log 2, 0)$, then extends back to $-\infty$ above the line $\text{Im } z = 0$. Since L is homotopic to the Hankel contour C defined in [5, 6], then by [6, Equation (25)], we have, for $s \in \mathbb{C} \setminus \mathbb{Z}$, the representation

$$\zeta_H(s) = \pi \cot(\pi s) \zeta(s) + \zeta(s+1) + \Gamma(1-s)\phi(s) - \psi(1-s)\zeta(s) - \zeta'(s), \quad (11)$$

where

$$\phi(s) := \frac{1}{2i\pi} \int_L \frac{z^{s-1} e^z}{e^z - 1} \text{Log}\left(\frac{e^z - 1}{z}\right) dz.$$

We now show that equations (9) and (11) are equivalent. To do this, we proceed as follows. Differentiation of the integral representation

$$\zeta(s) = \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1} e^z}{1 - e^z} dz$$

[5, Equation (2.4)], leads to the following identity:

$$\psi(1-s)\zeta(s) + \zeta'(s) = \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1} e^z \text{Log}(z)}{1 - e^z} dz \quad (s \neq 1, 2, 3, \dots) \quad (12)$$

[5, Equation (2.7)]. Splitting the right-hand member of (12), we can write

$$\frac{1}{2i\pi} \int_L \frac{z^{s-1} e^z \text{Log}(z)}{1 - e^z} dz = \phi(s) - \Phi(s), \quad (13)$$

with

$$\Phi(s) = \frac{1}{2i\pi} \int_L \frac{z^{s-1} e^z}{e^z - 1} \text{Log}(e^z - 1) dz.$$

From (12) and (13), we deduce the identity

$$\Gamma(1-s)\phi(s) - \psi(1-s)\zeta(s) - \zeta'(s) = \Gamma(1-s)\Phi(s).$$

Moreover, the substitution

$$z = \text{Log}(1 + e^{ix}), \quad -\pi < x < \pi,$$

in the contour integral defining Φ gives this function the equivalent form (10), proving that (9) and (11) are equivalent equations. \square

3.1 New formulas for $\gamma^{(k)}$

We deduce from (9) an evaluation of $\gamma^{(k)}$ which seems to us more meaningful than the one given in [15, Equation (21)].

Corollary 1. For any non-negative integer n , let \mathcal{L}_n and ξ_n be defined as follows:

$$\mathcal{L}_n := \Phi^{(n)}(1) = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} x \text{Log}^n(\text{Log}(1 + e^{ix})) dx,$$

and

$$\xi_n := (-1)^n \Gamma^{(n)}(1) = (-1)^n \int_0^{+\infty} e^{-x} \log^n(x) dx.$$

Then we have

$$\gamma^{(0)} = -\mathcal{L}_1 - \gamma_1 - \zeta(2), \quad (14)$$

and, for all positive integers k , the following general relations hold true:

$$\begin{aligned} \gamma^{(2k-1)} = & -\zeta^{(2k-1)}(2) - \frac{\gamma_{2k}}{2k} + \frac{1}{2k} \sum_{n=0}^{2k-1} \binom{2k}{n} \xi_n \mathcal{L}_{2k-n} \\ & + \sum_{n=1}^k \frac{2(2k-1)!}{(2k-2n)!} \gamma_{2k-2n} \zeta(2n), \quad (15) \end{aligned}$$

and

$$\begin{aligned} \gamma^{(2k)} = & \zeta^{(2k)}(2) - \frac{\gamma_{2k+1}}{2k+1} - \frac{1}{2k+1} \sum_{n=0}^{2k} \binom{2k+1}{n} \xi_n \mathcal{L}_{2k+1-n} \\ & + \sum_{n=1}^k \frac{2(2k)!}{(2k+1-2n)!} \gamma_{2k+1-2n} \zeta(2n) - 2(2k)! \zeta(2k+2). \end{aligned} \quad (16)$$

Proof of Corollary 1. The key formula to derive the general relations (15) and (16) is the representation of $\zeta_H(s)$ given by (9). Fortunately, the Laurent series expansion of each component in (9) can be written explicitly.

a) The Laurent expansions of $\pi \cot(\pi s)$ and $\zeta(s)$ at $s = 1$ are respectively

$$\pi \cot(\pi s) = \frac{1}{s-1} - \sum_{k \geq 1} 2\zeta(2k)(s-1)^{2k-1},$$

and

$$\zeta(s) = \frac{1}{s-1} + \gamma + \sum_{k \geq 1} \frac{(-1)^k}{k!} \gamma_k (s-1)^k,$$

where the coefficients γ_k are the Stieltjes constants. The expansion of $\pi \cot(\pi s) \zeta(s)$ is then obtained by Cauchy product:

$$\begin{aligned} \pi \cot(\pi s) \zeta(s) = & \frac{1}{(s-1)^2} + \frac{\gamma}{s-1} - \gamma_1 - 2\zeta(2) - \left(-\frac{1}{2}\gamma_2 + 2\gamma\zeta(2)\right)(s-1) \\ & + \frac{1}{2} \left(-\frac{1}{3}\gamma_3 + 4\zeta(2)\gamma_1 - 4\zeta(4)\right)(s-1)^2 \\ & - \frac{1}{6} \left(-\frac{1}{4}\gamma_4 + 6\zeta(2)\gamma_2 + 12\gamma\zeta(4)\right)(s-1)^3 + \dots \end{aligned} \quad (17)$$

b) It follows from the definition of ξ_k as $(-1)^k \Gamma^{(k)}(1)$ that the Laurent expansion of $\Gamma(1-s)$ at $s = 1$ is given by

$$\Gamma(1-s) = -\frac{1}{s-1} - \gamma - \sum_{k=2}^{\infty} \frac{\xi_k}{k!} (s-1)^{k-1} \quad (|s-1| < 1),$$

On the other hand, the function Φ defined by (10) is an entire function of s with $\Phi(1) = \mathcal{L}_0 = 0$, and the definition of \mathcal{L}_k as $\Phi^{(k)}(1)$ implies that the Taylor expansion of $\Phi(s)$ at $s = 1$ is given by

$$\Phi(s) = \sum_{k=1}^{\infty} \frac{\mathcal{L}_k}{k!} (s-1)^k.$$

The Laurent expansion of $\Gamma(1-s)\Phi(s)$ then follows by Cauchy product:

$$\begin{aligned}\Gamma(1-s)\Phi(s) &= -\mathcal{L}_1 - \left(\frac{1}{2}\mathcal{L}_2 + \gamma\mathcal{L}_1\right)(s-1) \\ &\quad + \frac{1}{2}\left(-\frac{1}{3}\mathcal{L}_3 - \gamma\mathcal{L}_2 - 2\gamma^{(0)}\mathcal{L}_1\right)(s-1)^2 \\ &\quad - \frac{1}{6}\left(\frac{1}{4}\mathcal{L}_4 + \gamma\mathcal{L}_3 + 3\gamma^{(0)}\mathcal{L}_2 + (2\zeta(3) + 3\gamma\zeta(2) + \gamma^3)\mathcal{L}_1\right)(s-1)^3 + \dots\end{aligned}\quad (18)$$

c) The Taylor expansion of $\zeta(s+1)$ at $s=1$ can be written as follows:

$$\zeta(s+1) = \zeta(2) + \sum_{k=1}^{\infty} \frac{(-1)^k}{k!} (-1)^k \zeta^{(k)}(2) (s-1)^k. \quad (19)$$

By assembling equations (17), (18), and (19) above, we obtain, by identifying the constant term, the relation (14). In the same way, the general formulae (15) and (16) are derived by identifying the coefficients of higher degree. \square

Remark 2. a) Combining (14) with the expression of $\gamma^{(0)}$ given by (1), we deduce the explicit evaluation:

$$\mathcal{L}_1 = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} x \operatorname{Log}(\operatorname{Log}(1+e^{ix})) dx = -\frac{3}{2}\zeta(2) - \frac{1}{2}\gamma^2 - \gamma_1. \quad (20)$$

b) The constants ξ_n involved in formulae (15)–(16) have a closed form expression in terms of Euler's constant γ and $\zeta(2), \dots, \zeta(n)$ [2, p. 227], [10, p. 12]. More precisely, we have $\xi_0 = 1$, $\xi_1 = \gamma$, and the recursion formula

$$\xi_{n+1} = \gamma\xi_n + \sum_{k=1}^n \frac{n!}{(n-k)!} \zeta(k+1) \xi_{n-k} \quad (n \geq 1),$$

so that

$$\begin{aligned}\xi_2 &= \gamma^2 + \zeta(2) = 2\gamma^{(0)}, \\ \xi_3 &= \gamma^3 + 3\gamma\zeta(2) + 2\zeta(3), \\ \xi_4 &= \gamma^4 + 6\gamma^2\zeta(2) + 8\gamma\zeta(3) + 3(\zeta(2))^2 + 6\zeta(4), \text{ etc.}\end{aligned}$$

A clever reformulation of this result is

$$\xi_n = P_n(\gamma, \zeta(2), \dots, \zeta(n)) \quad (n \geq 2),$$

where the polynomials P_n are defined by the exponential generating function

$$\exp\left(\sum_{k=1}^{\infty} x_k \frac{t^k}{k}\right) = \sum_{n=0}^{\infty} P_n(x_1, x_2, \dots, x_n) \frac{t^n}{n!}.$$

c) By mean of a theorem of Ramanujan [11, Theorem 4] whose proof is given in [4, p. 224], we can write the following identity:

$$(-1)^n \zeta^{(n)}(2) = n! + \gamma_n + \sum_{k=1}^{\infty} \frac{(-1)^k}{k!} \gamma_{k+n} \quad (n \geq 1).$$

Example 2. Applying formulas (15) and (16) to the case $k = 1$, and using (20), we obtain respectively

$$\gamma^{(1)} = -\zeta'(2) - \frac{1}{2}\gamma^3 - \frac{1}{2}\gamma_2 + \frac{1}{2}\mathcal{L}_2 + \frac{1}{2}\gamma\zeta(2) - \gamma\gamma_1,$$

and

$$\gamma^{(2)} = \zeta''(2) + \frac{1}{2}\gamma^4 - \frac{1}{3}\gamma_3 - \frac{1}{3}\mathcal{L}_3 - \gamma\mathcal{L}_2 + (2\gamma^2 + 5\gamma_1)\zeta(2) + \gamma^2\gamma_1 - \frac{1}{10}(\zeta(2))^2.$$

Numerical evaluations of the constants \mathcal{L}_2 and \mathcal{L}_3 are

$$\mathcal{L}_2 = -1.924491\dots \quad \text{and} \quad \mathcal{L}_3 = 7.158075\dots$$

3.2 New formulas for $\zeta(n)$ and $\zeta'(n)$

Corollary 2. For all integers $n \geq 2$, we have

$$\zeta(n) = \frac{(-1)^n}{(n-1)!} \times \frac{1}{2i\pi} \int_{-\pi}^{\pi} x \left(\text{Log}(1 + e^{ix})\right)^{n-1} dx, \quad (21)$$

and

$$\begin{aligned} \zeta'(n) = \frac{(-1)^n}{(n-1)!} \times \frac{1}{2i\pi} \int_{-\pi}^{\pi} x \text{Log} \left(\text{Log}(1 + e^{ix})\right) \left(\text{Log}(1 + e^{ix})\right)^{n-1} dx \\ + \zeta_H(n) - \zeta(n) \psi(n) - \zeta(n+1). \end{aligned} \quad (22)$$

Proof of Corollary 2. For $n \geq 2$, we can write

$$\begin{aligned} \pi \cot(\pi s) \zeta(s) &= \frac{\zeta(n)}{s-n} + \zeta'(n) + \text{O}(s-n), \\ \Gamma(1-s) &= \frac{(-1)^n}{(n-1)!} \left(\frac{1}{s-n} + (\gamma - H_{n-1}) \right) + \text{O}(s-n), \end{aligned}$$

and

$$\Gamma(1-s) \Phi(s) = \frac{(-1)^n}{(n-1)!} \frac{\Phi(n)}{s-n} + \frac{(-1)^n}{(n-1)!} (\Phi'(n) + \Phi(n)(\gamma - H_{n-1})) + \text{O}(s-n).$$

By applying formula (9) around $s = n$, we deduce the equation

$$\begin{aligned} \zeta_H(n) + O(s - n) &= \left(\zeta(n) + \frac{(-1)^n}{(n-1)!} \Phi(n) \right) \frac{1}{s-n} \\ &+ \frac{(-1)^n}{(n-1)!} (\Phi'(n) + \Phi(n)(\gamma - H_{n-1})) + \zeta'(n) + \zeta(n+1) + O(s-n). \end{aligned}$$

This leads to the identities

$$\zeta(n) + \frac{(-1)^n}{(n-1)!} \Phi(n) = 0,$$

and

$$\zeta_H(n) = \frac{(-1)^n}{(n-1)!} (\Phi'(n) + \Phi(n)(\gamma - H_{n-1})) + \zeta'(n) + \zeta(n+1),$$

thus proving formulas (21) and (22). \square

Corollary 3. For all positive integers n , we have

$$\Phi(2n+1) = (2n)! \zeta(2n+1), \quad (23)$$

and

$$-\Phi'(2n) = (2n-1)! (\zeta'(2n) + \zeta(2n) \psi(2n) - n\zeta(2n+1) + \sum_{k=1}^{n-1} \zeta(2n-k) \zeta(k+1)). \quad (24)$$

Proof of Corollary 3. By Euler's formula [4, p. 252], we have

$$\zeta_H(2n) = (n+1) \zeta(2n+1) - \sum_{k=1}^{n-1} \zeta(2n-k) \zeta(k+1),$$

and thus formulas (23) and (24) derive respectively from (21) and (22). \square

Example 3. Applying (23) and (24) for $n = 1$, we obtain

$$\Phi(3) = 2\zeta(3) = \zeta_H(2) = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} x \operatorname{Log}^2(1 + e^{ix}) dx,$$

and

$$-\Phi'(2) = \zeta'(2) + \zeta(2)(1-\gamma) - \zeta(3) = \frac{1}{2i\pi} \int_{-\pi}^{\pi} x \operatorname{Log}(1 + e^{ix}) \operatorname{Log}(\operatorname{Log}(1 + e^{ix})) dx.$$

Remark 3. It should be noted that our formula (21) for $\zeta(n)$ is similar to but different from that given in [5, Equation (2.6)] which, after substitution $z = \operatorname{Log}(1 + e^{ix})$, translates into

$$\zeta(n) = \frac{(-1)^{n-1}}{(n-1)!} \times \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Log}(\operatorname{Log}(1 + e^{ix})) (\operatorname{Log}(1 + e^{ix}))^{n-1} dx.$$

3.3 New formulas for Glasser-Manna-Oloa type integrals

Using the relation between $\Phi(s)$ and $\zeta_H(s)$ arising from formula (9) in a neighborhood of $s = -n$, we derive new formulas for Glasser-Manna-Oloa type integrals.

Corollary 4. We have

$$\Phi(0) = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} \frac{x}{\operatorname{Log}(1 + e^{ix})} dx = \frac{1}{2} \log(2\pi) - \frac{1}{2}\gamma + \frac{1}{2}, \quad (25)$$

and

$$\begin{aligned} \Phi'(0) &= -\frac{1}{2i\pi} \int_{-\pi}^{\pi} \frac{x \operatorname{Log}(\operatorname{Log}(1 + e^{ix}))}{\operatorname{Log}(1 + e^{ix})} dx \\ &= \frac{1}{2}\gamma_1 + \frac{1}{4} \log^2(2\pi) + \frac{1}{4}\gamma^2 - \frac{1}{2}\gamma - \frac{1}{2}\gamma \log(2\pi) - \frac{7}{8}\zeta(2) + \beta, \end{aligned} \quad (26)$$

where β is the linear coefficient in the Laurent expansion of ζ_H at $s = 0$.

Proof of Corollary 4. Around 0, we have the decomposition given by (9):

$$\zeta_H(s) = \zeta(s + 1) + \pi \cot(\pi s) \zeta(s) + \Gamma(1 - s)\Phi(s),$$

and the expansion [9, Equation (8)]

$$\zeta_H(s) = \frac{1}{2s} + \frac{1}{2}\gamma + \frac{1}{2} + \beta s + O(s^2).$$

On the other side, we have the expansions

$$\begin{aligned} \zeta(s + 1) &= \frac{1}{s} + \gamma - \gamma_1 s + O(s^2), \\ \pi \cot(\pi s) \zeta(s) &= -\frac{1}{2s} - \frac{1}{2} \log(2\pi) + \left(\frac{1}{2}\gamma_1 - \frac{1}{4} \log^2(2\pi) + \frac{1}{4}\gamma^2 + \frac{7}{8}\zeta(2) \right) s + O(s^2), \\ \Gamma(1 - s)\Phi(s) &= \Phi(0) + (\Phi'(0) + \gamma\Phi(0)) + O(s^2). \end{aligned}$$

By identifying the constant coefficient in the expansion of the right-hand member of (9), we deduce the equation

$$\frac{1}{2}\gamma + \frac{1}{2} = \gamma - \frac{1}{2} \log(2\pi) + \Phi(0),$$

which is equivalent to (25). In the same way, formula (26) is derived by identifying the linear coefficient in the expansion of the right-hand member of (9). \square

Remark 4. We derive from [17, Equation (53)] the following evaluation of the constant β which appears in (26):

$$\beta = -\gamma_1^{[2]}(0) - \gamma_1 = 1 + \frac{1}{2}\gamma - \frac{1}{4}\gamma^2 - \gamma_1 + \frac{1}{4}\zeta(2) - \sum_{n=2}^{\infty} \frac{|b_n|}{(n-1)^2} = 1.589935\dots,$$

where b_n are the Bernoulli numbers of the second kind defined by way of their generating function

$$\frac{x}{\log(x+1)} = \sum_{n=0}^{\infty} b_n x^n = 1 + \frac{x}{2} - \frac{x^2}{12} + \frac{x^3}{24} - \frac{19x^4}{720} + \frac{3x^5}{160} - \frac{863x^6}{60480} + \dots$$

Remark 5. Formula (25) appears in a slightly different but equivalent form in [14, Equation (3)], due to the identities

$$\begin{aligned} \Phi(0) &= \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{iy \, dy}{2 \log(2 \cos(\frac{y}{2})) + iy} \\ &= \frac{2}{\pi} \int_0^{\pi} \frac{y^2}{y^2 + 4 \log^2(2 \cos(\frac{y}{2}))} \, dy \\ &= \frac{4}{\pi} \int_0^{\frac{\pi}{2}} \frac{x^2}{x^2 + \log^2(2 \cos(x))} \, dx. \end{aligned}$$

Corollary 5. We have

$$\Phi(-1) = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} \frac{x}{\text{Log}^2(1 + e^{ix})} \, dx = \log(\mathcal{A}) - \frac{1}{12}\gamma + \frac{7}{24}, \quad (27)$$

where $\mathcal{A} = \exp(\frac{1}{12} - \zeta'(-1))$ is the Glaisher-Kinkelin constant, and for all integers $n \geq 2$,

$$(2n-1)! \Phi(1-2n) = \log(\mathcal{A}_n) - \frac{B_{2n}}{2n}\gamma + \frac{1}{2n} \sum_{k=0}^{2n-2} \binom{2n}{k} \frac{B_k B_{2n-k}}{2n-k}, \quad (28)$$

with $\log(\mathcal{A}_n) := \frac{H_{2n-1} B_{2n}}{2n} - \zeta'(1-2n)$.¹

Proof of Corollary 5. Around $s = -1$, we have the expansion [9, Equation (9)]

$$\zeta_H(s) = -\frac{1}{12(s+1)} - \frac{1}{12}\gamma - \frac{1}{8} + O(s+1).$$

¹The numbers \mathcal{A}_n are called generalized Glaisher-Kinkelin constants [12].

On the other side, we have

$$\begin{aligned}\zeta(s+1) &= -\frac{1}{2} + O(s+1), \\ \pi \cot(\pi s) \zeta(s) &= \zeta'(-1) + O(s+1), \\ \Gamma(1-s)\Phi(s) &= \Phi(-1) + O(s+1).\end{aligned}$$

Formula (27) is then deduced by identifying the constant coefficient in the expansion of the right-hand member of (9). In the same way, from [9, Proposition 10], we have for $n \geq 2$ the Laurent expansion around $s = 1 - 2n$:

$$\zeta_H(s) = \frac{\zeta(1-2n)}{s+2n-1} - \frac{B_{2n}}{2n}\gamma + \frac{H_{2n-1}B_{2n}}{2n} + \frac{1}{2n} \sum_{k=0}^{2n-2} \binom{2n}{k} \frac{B_k B_{2n-k}}{2n-k} + O(s+2n-1)$$

which, by the same method, allows us to deduce formula (28). \square

Corollary 6. For all positive integers n , we have the identity

$$\Phi(-2n) = \frac{(-1)^{n+1}}{2} (2\pi)^{-2n} \zeta(2n+1) + \frac{2n+1}{4n} \frac{B_{2n}}{(2n)!}, \quad (29)$$

from which can be deduced the following relation:

$$2(2n)! \Phi(-2n) = (-1)^{n+1} (2\pi)^{-2n} \Phi(2n+1) + \frac{2n+1}{2n} B_{2n}. \quad (30)$$

Proof of Corollary 6. Around $s = -2n$, we have

$$\zeta_H(s) = -\frac{B_{2n}}{4n} + \frac{B_{2n}}{2} + O(s+2n).$$

On the other side, we have the expansions

$$\begin{aligned}\zeta(s+1) &= -\frac{B_{2n}}{2n} + O(s+2n), \\ \pi \cot(\pi s) \zeta(s) &= \zeta'(-2n) + O(s+2n), \\ \Gamma(1-s)\Phi(s) &= (2n)! \Phi(-2n) + O(s+2n).\end{aligned}$$

By identifying the constant coefficient in the expansion of the right-hand member of (9), we obtain

$$\begin{aligned}(2n)! \Phi(-2n) &= -\zeta'(-2n) + \frac{2n+1}{4n} B_{2n} \\ &= \frac{(-1)^{n+1}}{2} (2\pi)^{-2n} (2n)! \zeta(2n+1) + \frac{2n+1}{4n} B_{2n},\end{aligned}$$

which is nothing else than (29). Moreover, using the identity

$$\Phi(2n+1) = (2n)! \zeta(2n+1)$$

arising from (23) allows us to deduce the relation (30). \square

References

- [1] T. Apostol and T. H. Vu, Dirichlet series related to the Riemann zeta function, *J. Number Theory* **19** (1984), 85–102.
- [2] T. Apostol, Formulas for higher derivatives of the Riemann zeta function, *Math. of Computation*, **44** (1985), 223–232.
- [3] U. Balakrishnan, Laurent Expansion of Dirichlet Series-II, *Monatshefte für Mathematik*, **103** (1987), 1–5.
- [4] B. C. Berndt, *Ramanujan's Notebooks Part I*, Springer Verlag, New York, 1985.
- [5] K.N. Boyadzhiev, Evaluation of series with Hurwitz and Lerch zeta function coefficients by using Hankel contour integrals, *Applied Mathematics and Computation* **186** (2007), 1559– 1571.
- [6] K.N. Boyadzhiev, H. Gadiyar, and R. Padma, The values of an Euler sum at the negative integers and a relation to a certain convolution of Bernoulli numbers, *Bulletin of the Korean Mathematical Society* **45** (2008), 277–283.
- [7] W. E. Briggs and R. G. Buschman, The power series coefficients of functions defined by Dirichlet series, *Illinois J. Math.* **5** (1961), 43–44.
- [8] B. Candelpergher, *Ramanujan Summation of Divergent Series*, Lecture Notes in Math. 2185, Springer, 2017.
- [9] B. Candelpergher and M-A. Coppo, Laurent expansion of harmonic zeta functions, *Journal of Mathematical Analysis and Applications* **491** (2020), 124309.
- [10] Junesang Choi and H. M. Srivastava, Evaluation of higher-order derivatives of the Gamma function, *Publikacije Elektrotehnickog fakulteta. Serija Matematika* **11** (2000), 9–18.
- [11] B. K. Choudhury, The Riemann Zeta-Function and Its Derivatives, *Proceedings Royal Society London* **450** (1995), 477–499.
- [12] M-A. Coppo, Generalized Glaisher-Kinkelin constants and Ramanujan summation of series, *Research in Number Theory* **10**, 15 (2024).
- [13] M-A. Coppo and P. T. Young, On shifted Mascheroni series and hyperharmonic numbers, *J. Number Theory*, **169** (2016), 1–20.

- [14] M. L. Glasser and D. Manna, *On the Laplace transform of the psi-function*, Tapas in Experimental Mathematics (T. Amdeberhan and V. Moll, eds.), Contemporary Mathematics, vol. 457, American Mathematical Society, 2008, 193–202.
- [15] L. Kargin, A. Dil, M. Cenkci, and M. Can, On the Stieltjes constants with respect to harmonic zeta functions *Journal of Mathematical Analysis and Applications* **525** (2023), 127302.
- [16] O. Oloa, *Some Euler-type integrals and a new rational series for Euler's constant*, Tapas in Experimental Mathematics (T. Amdeberhan and V. Moll, eds.), Contemporary Mathematics, vol. 457, American Mathematical Society, 2008, 237–248.
- [17] P. T. Young, Global series for height 1 multiple zeta functions, *European Journal of Mathematics* **9** (2023), 9:99.